Athlete Name:	Date of Concussion Diagnosis:
---------------	-------------------------------

MUHSD Concussion Return to Play (RTP) Protocol

CA STATE LAW AB 2127 (Effective 1/1/15) STATES THAT RETURN TO PLAY (I.E., COMPETITION) CANNOT BE SOONER THAN 7 DAYS AFTER EVALUATION BY A PHYSICIAN (MD/DO/PA/NP) WHO HAS MADE THE DIAGNOSIS OF CONCUSSION.

Instructions:

- This graduated return to play protocol MUST be completed before you can return to FULL COMPETITION.
 - The athletic trainer (AT), physician, and/or identified concussion monitor (e.g., Athletic Director, Head Coach under supervision of athletic director), must monitor your progression and initial each stage after you successfully pass it.
 - Stages I to II-D take a minimum of 6 days to complete.
 - You must complete one full practice without restrictions (Stage III) before competing in first game.
- After Stage I, you cannot progress more than one stage per day (or longer if instructed by your physician).
- If symptoms return at any stage in the progression, IMMEDIATELY STOP any physical activity and follow up with your schools AT, other
 identified concussion monitor, or your physician. In general, if you are symptom-free the next day, return to the previous stage where
 symptoms had not occurred.
- Seek further medical attention if you cannot pass a stage after 3 attempts due to concussion symptoms, or if you feel uncomfortable at any time during the progression.

Stage	Activity	Exercise Example	Objective of the Stage	Date & Initials (A.D., A.T. or Head Coach)
ı	No physical activity for at least 2 full symptom-free days AFTER you have seen a physician	No activities requiring exertion (weight lifting, jogging, P.E. classes)	Recovery and elimination of symptoms	
II-A	Light aerobic activity	 10-15 minutes (min) of walking or stationary biking. Must be performed under direct supervision by designated individual 	 Increase heart rate to no more than 50% of perceived maximum (max) exertion (e.g., 100 beats per min) Monitor for symptom return 	
II-B	Moderate aerobic activity (Light resistance training)	 20-30 min jogging or stationary biking Body weight exercises (squats, planks, push- ups), max 1 set of 10, no more than 10 min total 	 Increase heart rate to 50-75% max exertion (e.g.,100-150 bpm) Monitor for symptom return 	
II-C	Strenuous aerobic activity (Moderate resistance training)	 30–45 min running or stationary biking Weight lifting ≤ 50% of max weight 	 Increase heart rate to > 75% max exertion Monitor for symptom return 	
II-D	Non-contact training with sport-specific drills (No restrictions for weightlifting)	 Non-contact drills, sport-specific activities(cutting, jumping, sprinting) No contact with people, padding or the floor/mat 	 Add total body movement Monitor for symptom return 	
Minim	um of 6 days to pass Stages I and play, after successful co	II. Prior to beginning Stage III, please make sure ompletion of Stages I and II, has been given to yo	that written physician (MD/DO) clearance ur school's concussion monitor	for return to
III	Limited contact practice	Controlled contact drills allowed (no scrimmaging)	 Increase acceleration, deceleration and rotational forces Restore confidence, assess 	
	Full contact practice Full unrestricted practice	Return to normal training, with contact Return to normal unrestricted training	readiness for return to play Monitor for symptom return	
MA		t least ONE contact practice before return to com		cted practice
IV	Return to play (competition)	Normal game play (competitive event)	Return to full sports activity without restrictions	